

Isolation and determination of deoxynivalenol by reversed-phase high-pressure liquid chromatography

Abstract

Deoxynivalenol (DON) is a mycotoxin produced by food contamination. It is a pharmacologically active compound that acts on the serotonin receptor, leading to several neuroendocrine and hematological disorders. In this article we describe a simple, accurate, and sensitive method for the quantification of DON. DON was quantified using a Phenomenex® ODS analytical C18 column (150 mm × 46 mm, 5 µm) with a mobile phase composed of mixture of water-acetonitrile-methanol (5:4:1, v/v/v) at a flow rate of 1.5 ml/min and at 254 nm in an ultraviolet (UV) detector. The method has been validated with isolated samples of DON and provides a tool for the control of substandard and counterfeit commercial food products.

Key words: Deoxynivalenol; *Fusarium graminearum*; HPLC; mycotoxins

Vikas Kumar Gupta,
Pronobesh Chattopadhyay,
Mohan Ch Kalita¹, Asshwani
Kumar Chaurasia, Hemanta
Kumar Gogoi, Lokendra Singh

Division of Biotechnology,
Defense Research Laboratory,
Post Bag No: 2, Tezpur-784 001,
¹Gauhati University, Department
of Biotechnology, Guwahati-781
014, Assam, India

Address for correspondence:

Dr. Pronobesh Chattopadhyay
Defense Research Laboratory,
Post Bag No: 2, Tezpur-784 001,
Assam, India.
Email-chattopadhyay.drdo@gmail.
com

Access this article online

Website: www.phmethods.org

DOI: 10.4103/2229-4708.81087

Quick response code



INTRODUCTION

Mycotoxins, which are by-products of fungal metabolism, can cause adverse health reactions in humans and animals consuming infected agricultural products.^[1,2] Mycotoxins are usually trichothecenes, which have low molecular weight, are lipophilic in nature, and accumulate in the fat fractions of plants. They are toxic to animals.^[3] Trichothecenes are a diverse range of structurally related compounds derived from 12,13-epoxytrichothec-9-ene. Two groups can be distinguished according to the chemical structure at carbon-8 (C-8): group A trichothecenes have a functional group other than a ketone at C-8 and group B trichothecenes are characterized by a ketone at C-8.^[4] Deoxynivalenol (DON), also known as vomitoxin (VT) and 12,13-epoxy-3,7,15-trihydroxy-trichothec-9-en-8-one [Figure 1]), is a trichothecene mycotoxin produced by *Fusarium graminearum* and *F. culmorum* and a frequent cause of toxicity after consumption of grain-based agricultural products,^[5] being responsible for a variety of mycotoxicoses in animals and humans.^[6]

DON is usually separated by an internal-surface reversed-phase column (octadecyl silica column) and analyzed by liquid chromatography–mass spectrometry (LC–MS), using post-derivatization with 1-anthroylnitrile.^[7] Gas chromatography–mass spectrometry (GC–MS) and GC–MSMS has been used to detect verrucarol (VER) and trichodermol (TRID) but so far there have been few reports of the use of high-pressure liquid chromatography (HPLC) and ultraviolet (UV) detector for analysis of DON. HPLC methods are more reliable and more accurate than other analytical techniques. RIA and enzyme-linked immunosorbent assay (ELISA)^[8] are two methods that are most commonly used for DON analysis, but both involve multiple steps and are complicated time-consuming procedures and, besides, the results are sometimes non-reproducible. Therefore, the present study was designed to develop a methodology for estimation of DON by reversed-phase high-pressure liquid chromatography (RPHPLC) using a C-18 column and a UV detector.




Figure 1: Structure of deoxynivalenol

MATERIALS AND METHODS

Materials

Four commercially available cereals, viz, corn, wheat, rice, and *kabli matar* (peas) were purchased from a randomly selected local vendor in Tezpur, Assam, India. Standard DON was purchased from Sigma-Aldrich (Madrid, Spain) [batch No: 303, filling code: 1364166, purity: $\geq 97.0\%$ (TLC)]. The organic solvents used were HPLC grade and were purchased from Waters® (Milford, MA, USA). The other chemicals, charcoal, alumina, and silica (60–120 mesh), were obtained from Merck® (Darmstadt, Germany).

Isolates

MTCC strain 1893 *Fusarium graminearum* was obtained from the Institute of Microbial Technology (IMTECH) Chandigarh, India, and incubated as described below.

Incubation

Four conical flasks (250 ml) containing dry polished rice, corn, wheat, and peas (*kabli matar*) (100 g) were moistened with 30 ml deionized water and kept overnight in a controlled atmosphere having suitable equilibrium relative humidity (RH) (temp $25 \pm 5^\circ\text{C}$ and RH 70%). The cereals were then autoclaved at 120°C for 30 min and the flasks were inoculated with 5 ml spore suspension containing approximately 10^6 macroconidia. The suspension was made from a single spore culture. The cultures were incubated (temp $20 \pm 8^\circ\text{C}$ and RH $70 \pm 5\%$) for 3 weeks for fermentation. Five-gram samples were finely ground in a laboratory mill and thoroughly mixed before aliquots were taken for the experiment.

Extraction and cleanup

At the end of the incubation period the flasks were dried at $40 \pm 2^\circ\text{C}$ in a hot-air oven for 24 h. All samples were then dry ground to the consistency of flour in a

laboratory mill. Five-gram samples of ground grain were extracted with 30 ml of the extraction solvent (water:acetonitrile:methanol in a ratio of 5:4:1, v/v) in Erlenmeyer flasks, with continuous shaking for 30 min. The extraction solvent was filtered through Whatman™ No. 4 filter paper. For the cleanup process about 30 ml of each extract was placed in a 18×85 mm tube and filtered through a self-designed alumina-silica-activated charcoal (1:1:1) column and eluted in clear tips. The elute was evaporated to dryness on a rotary evaporator. The residue was dissolved in 1 ml of water:acetonitrile:methanol (5:4:1). Each extract was further purified by filtration through a separate Whatman™ filter (Puradisc™ 25AS, $0.45\text{-}\mu\text{m}$ pore size) and an aliquot was used for the HPLC analysis.

Instruments

A Waters® 600E system (Waters®, Milford, MA, USA) provided with a programmable UV detector was used in this study. The injector system was Rheodyne® (USA), with a maximum capacity of $20\ \mu\text{l}$ and with Empower™ software. A reverse phase Phenomenex® ODS C18 column ($150\ \text{mm} \times 46\ \text{mm}$, $5\ \mu\text{m}$) protected with a guard was used throughout the experiments. Chromatograms were recorded and integrated using a Waters® 745 integrator. The UV monitor was set to 254 nm, with a sensitivity of 002 or 005 absorbance units full scale. The recorder was set to 10 min ($25\ \text{cm/h}$).

Validation studies

The rectilinear relationship between concentrations of the analytes and the UV detector response were evaluated. The concentrations used were 10, 20, 30, 40, 50, 60, 70, 80, 90, and 100 ng/ml for DON. Three different preparations of the analytical standard were analyzed in triplicate on the same day for determination of intra-day assay precision. These determinations were repeated using freshly prepared standard solutions on three separate days to determine inter-day precision of analysis.

Analytical solutions and isolated sample in triplicate, freshly prepared solutions on three separate days, were used to compute the inter-day ($n=6$ separate determinations) and intra-day precision of the method. The stability of the analytical solutions was determined for DON at the concentrations described above for the assessment of repeatability. Analytical solutions were injected repeatedly ($n=36$) and RSDs % computed for the peak areas due to the respective analytes.

The optimized method for recovery was applied to the

analysis of DON content in cereal samples by spiking of DON in wheat sample (level of contamination: 1 mg DON/kg).

The performance characteristics of the method were based on the resolution between the standard DON and isolated DON and the robustness of the method as a function of small changes in the ratio of the solvent (water:acetonitrile:methanol 6:3:1 v/v) of the mobile phase, stability of analytical solutions, variation of analyst, and the effect of temperature (20–40°C) on resolution. The limits of detection and quantization for each analytes were determined as S/N of three and RSD % $\leq 5\%$, respectively.

RESULTS

Increase in the water:acetonitrile ratio resulted in an improvement in the chromatographic peak shapes. But adding small amount of methanol increased the resolution of separation of DON. The optimum ratio was water:acetonitrile:methanol of 5:4:1 (v/v/v) and the most sharp and prominent peaks were observed at this ratio. This observation may have been due to competition between the highly basic moieties of DON and the hydronium ions from the methanol and water molecules for electrostatic interaction with residual amine [Figure 2]. Further, in keeping view that instability of some silica-based columns at pH values of below 28, the method is novel that all methods are using organic solvents.

Assessment of the effect of temperature (over a range of 18–40°C) on the separation of the analyte showed that there was a gradual, albeit insignificant, decrease in retention times with increase of temperature, which did not compromise the resolution of the separation of DON. Subsequently all analyses were performed at the ambient temperature ($\approx 20^\circ\text{C}$). The stability of analytical solutions of DON was assessed at three different concentrations (as described in the 'validation' section) and produced RSD % values ($n=36$ injections over 72 h) of peak areas which were typically less than 25%, 15%, and 05% at low, intermediate, and high concentrations, respectively. Thus, the solutions are sufficiently stable to justify analysis being done after preparation of fresh samples over 24-h periods. The chromatographic peak areas showed a rectilinear relationship to the concentration of the analyte within the specified ranges [Table 1] which is consistent with the expected concentrations on dilution of the innovator product of DON and isolated products. Linear regression analysis showed that the

correlation coefficients (r^2) of all calibration curves were ≥ 0.997 , with minimal variation in the slopes and intercepts [Table 2]. The performance characteristics and validation data for the method using the mobile water:acetonitrile:methanol are summarized in Table 3. The intra-day assay precision (RSD%) of peak areas for DON was limiting.

The values of recovery of DON concentrate obtained from six replicates of two samples at confidence interval at 97% probability which is accepted limit [Table 4].

DISCUSSIONS

The challenge involved in the measurement of DON in food is that the conventional methods are inaccurate.^[8,9] ELISA is the established method for




Figure 2: Chromatogram of DON toxin

Table 1: Regression analyses of calibration curves generated from the analysis of deoxynivalenol

Analyte	Range (ng/ μl)	n	Slope	Intercept correlation	Coefficient (R2)
DON	10 -80	6	40.161 \pm 0.1	63.11 \pm 4.2	0.9977 \pm 0.0001

Table 2: Repeatability and sensitivity of HPLC method

	Deoxynivalenol		
	Conc.	AUC	R.S.D (%)
Inter-day precision*			
Day 1 (n= 3)	12.5ng/ml,	76321 \pm 23.99,	1.11%
Day 2 (n=3)	25ng/ml,	221737 \pm 26.55,	1.65%
Intra-day precision*			
Solution 1(n=3)	12.5ng/ml,	75220 \pm 14.55,	0.561%
Solution 2(n=3)	25ng/ μg /ml,	221737 \pm 27.84,	0.73%
Limit of detection			
S/N (1mL)	0.04		
Limit of quantification*			
R.S.D. ≤ 5 (g/mL)	0.087		

*All precision data are mean \pm S.D. (n = 3) with R.S.D. % values in parenthesis. Precision data represents peak areas for analytes corrected for quantities of analytical standards materials used in the preparation of solutions. The concentrations in ng/ml represents the levels at which precision have been assessed. AUC: Area under Curve.

Table 3: Precision of RP-HPLC analysis of deoxynivalenol substances in isolated sample

		Deoxynivalenol
Inter-day precision*		
Mean \pm S.D. (n = 3) (% w/w)		80.12% \pm 0.66
R.S.D. %		0.70%
Intra-day precision*		
Mean \pm S.D. (n = 3) (% w/w)		82.63% \pm 0.58
R.S.D. %		0.83%

*All precision data are mean \pm S.D. (n = 3) with R.S.D. % values in parenthesis. Precision data represents peak areas for analytes corrected for quantities of analytical standards materials used in the preparation of solutions.

Table 4: Recoveries of deoxynivalenol for wheat and rice determined by HPLC method

Actual spiking deoxynivalenol		0.1 mg/kg	1 mg/kg
Concentration			
Analyst 1*	Wheat	87.90 \pm 4.50	89.67 \pm 4.55
	Rice	86.11 \pm 6.11	93.57 \pm 6.09
Analyst 2	Wheat	86.82 \pm 6.11*	87.99 \pm 5.90*
	Rice	86.04 \pm 5.22*	94.02 \pm 5.92**

All precision data are mean \pm S.D. (n = 6) Significantly different ($P < 0.05$, ** $P < 0.01$) from sham operated control group.

measurement of DON. DON is readily oxidized to the respective quinines, which are substances containing the highly basic quinoids moiety. This interacts with residual silanols on silica-based columns.^[10,11]

Pascale^[7] was the first to determine T₂ toxin (which are similar to DON in chemical structure) by liquid chromatography, with fluorescence detection derivatization with 1-anthroylnitrile. Though, after derivatization the molecules became stabilized for HPLC analysis, the accuracy and sensitivity of this method were questionable.

Determinations of DON by ELISA methods are rapidly using but RPHPLC by using UV detector rapid and simple methods are unavailable. Acetyl (CH₃C=O) is responsible for the toxicity of DON: 3-acetyl-DON is more prevalent in Europe and 15 acetyl-DON is more prevalent in North America.^[9,12] These toxins are often present at levels of 10%–20% in DON, and differ only with an acetyl group. The present investigation shows that using of CH₃OH in mobile phases provides a competing ion to reduce analyte–silanol interactions. The validation of the repeatability (inter- and intra-day precision) of the proposed HPLC method using the isolated product [Table 3] yielded RSD % values typically less than 2%. It is obvious that the difficulties in its synthesis, the absence of regulatory standards, and toxin counterfeiting have resulted in the marketing of products.

Compared with the other reported methods for estimation of DON, this method is simple and quick. A previous study showed that for isolation and increases maximum absorbance in UV range DON was passes through immunoaffinity column,^[13] whereas in the present study DON was separated using alumina-silica-activated charcoal (1:1:1), which is more economic and easier.

Previously, DON was isolated by Sephadex™ LH20 column^[14] and other methods, viz, solid phase extraction, liquid extraction, ion-exchange columns, immunoaffinity columns, etc.^[15] The new procedure described here for isolation of DON has several advantages: it is economical, simple, and less time consuming than the previous methods for large-scale separation of DON.

HPLC methods are sensitive and more accurate compared with LC–MS and GC–MS and most manufacturers prefer the HPLC methods.^[16] Besides, LC–MS and GC–MS are expensive techniques, and not all manufacturers are able to bear the costs involved. The present investigation presents an alternative to LC–MS for analysis for DON. The method has a novel approach, using an immunoaffinity column for isolation of DON.

CONCLUSIONS

DON is a potential toxin found in food grains. HPLC methods can be useful techniques for quality control and monitoring of this contamination. In this article we have described an accurate, sensitive, and easy technique to estimate DON in a food sample. The present method validation and repeatability compiles with standard. Therefore, the present investigation has importance for alternates of ELISA, LC-MS analysis for DON.

ACKNOWLEDGMENTS

This research was fully supported by the Defense Research Laboratory, DRDO, who provided all facilities and financial assistance.

REFERENCES

- Morris CM, Li YC, Ledoux DR, Bermudez AJ, Rottinghaus GE. The individual and combined effects of feeding moniliformin supplied by *Fusarium fujikuroi* culture material and deoxynivalenol in young turkey poults. *Poult Sci* 1999;78:1110-5.
- Müller G, Kielstein P, Rosner H, Berndt A, Heller M, Köhler H. Studies on the influence of combined administration of ochratoxin A

- fumonisin B1 deoxynivalenol and T-2 toxin on immune and actions in weaner pigs. *Mycoses* 1999;42:485-93.
3. Pronk MEJ, Schothorst RC, Van Egmond HP. Toxicology and occurrence of nivalenol fusarenon X diacetoxyscirpenol neosolaniol and 3-and 15-acetyldeoxynivalenol a review of six trichothecenes. *RIVM Rep* 2002;38880:2024.
 4. Cusumano V. Trichothecens - chemical biological and toxicological aspects. In: Ueno Y, editor. Chapter 4: Immunobiological activities of mould products: Functional impairment of human monocytes exposed to aflatoxin B₁. Netherland: Elsevier; 1983. p. 54.
 5. Sardjono NA, Yasmashita A, Yoshizawa T. Natural co-occurrence of aflatoxins and *Fusarium* mycotoxins (fumonisins deoxynivalenol nivalenol and zearalenone) in corn from Indonesia. *Food Addit Contam* 1998;15:337-84.
 6. Thiel PG. A molecular mechanism for the toxic action of moniliformin a mycotoxin produced by *Fusarium moniliforme*. *Biochem Pharmacol* 1978;27:483-9.
 7. Pascale M, Haidukowski M, Viscont A. Determination of T-2 toxin in cereal grains by liquid chromatography with fluorescence detection after immunoaffinitycolumn clean-up and derivatization with 1-anthroylnitrile. *J Chromatogr A* 2003;989:257-64.
 8. Trucksess MW, Thomas F, Young K, Stack ME, Fulgueras WJ, Page SW. Survey of deoxynivalenol in US 1993 wheat and barley crops by enzyme-linked immunosorbent assay. *J AOAC Int* 1995;78:631-6.
 9. Rotter BA, Prelusky DB, Pestka JJ. Toxicology of deoxynivalenol (vomitoxin). *J Toxicol Environ Health* 1996;48:1-34.
 10. Pestka JJ, Lin WS, Miller ER. Emetic activity of the trichothecene 15-acetyldeoxynivalenol in swine. *Food Chem Toxicol* 1987;25:855-8.
 11. Ehling G, Cockburn A, Snowdon P, Buchhaus H. The significance of the *Fusarium* toxin deoxynivalenol (DON) for human and animal health. *Cereal Res Commun* 1997;25:433-47.
 12. Riley RT. Mechanistic interactions of mycotoxins: Theoretical consideration. In: Sinha KK, Bhatanagar D, editor. Chapter 9. Mycotoxins in Agriculture and Food Safety. New York: Marcel Dekker; 1998. p. 227-54.
 13. Kotal F, Zazuna R. A simple method for determination of deoxynivalenol in cereals and flours. *Czech J Food Sci* 2002;20:63-8.
 14. Kenneth C, Ehrlich H, Eivind B, Lilleho J. Simple Method for Isolation of 4-Deoxynivalenol from Rice Inoculated with *Fusarium graminearum*. *Appl Environ Microbiol* 1984;48:1053-4.
 15. Rahmani A, Jinap S, Soleimany F. Qualitative and quantitative analysis of mycotoxin. *Comp Rev Food Sci Food Safety* 2009;8:202-51.
 16. Qiu P, Chen X, Chen L, Lin C. Simultaneously determination of five alkaloids in body fluids by high performance liquid chromatography coupled with electro spray ionization tandem mass spectroscopy. *J Chromatogr B* 2008;875:471-7.

Source of Support: Defense Research Laboratory, DRDO,
Conflict of Interest: None declared.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) **First Page File:**

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) **Article File:**

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1024 kb. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) **Images:**

Submit good quality color images. Each image should be less than **4096 kb (4 MB)** in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) **Legends:**

Legends for the figures/images should be included at the end of the article file.